

BEYOND BFKL

R. B. PESCHANSKI

*CEA/DSM/SPhT, Unité de recherche associée au CNRS,
CE-Saclay, F-91191 Gif-sur-Yvette Cedex, France;
Email: pesch@spht.saclay.cea.fr*

The Balitsky-Fadin-Kuraev-Lipatov (BFKL) evolution equation is known to be “unstable” with respect to fluctuations in gluon virtuality, transverse momentum and energy requiring to go beyond the leading order BFKL. Still, these instabilities point to fruitful improvements of our deep understanding of QCD. Recent applications to next-leading order and to saturation problems are outlined.

1. “Instabilities” of the BFKL Equation




Figure 1. *The QCD evolution landscape.* The BFKL evolution and its limitations. **i)** to the left: non-perturbative region; **ii)** to the right: DGLAP evolution; **iii)** to the top: Saturation region.

The Balitsky-Fadin-Kuraev-Lipatov (BFKL) evolution equation has already a venerable past¹. It appears as a key tool in many recent works on small-x physics (in the broad sense). It is interesting to notice that its limitations themselves are the seeds of interesting fields of research. Let us discuss limitations which can be associated with the idea of “instabilities”.

- **i)** *Instability towards the non-perturbative regime*

It is well known that the perturbative “gluon ladder” contributing to the BFKL cross-section is characterized by a “cigar-shape” structure² of the transverse momenta. Hence, it is difficult to avoid an excursion inside the near-by non-perturbative region

(Fig. (1), to the left).

- **ii)** *Instability towards the renormalization group regime*

Calculations of the next-leading BFKL kernel³ has proven that the inclusion of next-leading logs gives a (too) strong correction to the leading log result. After a resummation motivated by the suppression of spurious singularities^{4,5}, the results show that the resummed NLO-BFKL kernels are very similar, *e.g.* “attracted” towards the Dokshitzer Gribov Lipatov Altarelli Parisi (DGLAP) evolution⁶, at least for the structure functions

(Fig. (1), to the right).

- **iii)** *Instability towards the high density (saturation) regime*

The BFKL evolution implies a densification of gluons and sea quarks, while they keep in average the same size. It is thus natural to expect⁷ a modification of the evolution equation by non-linear contributions in the gluon density. Recently, the corresponding theoretical framework has been settled^{8,9}, and is based on an extension of the BFKL kernel acting on non-linear terms. It leads to a transition to the saturation regime (Fig. (1), to the top).

The main subjects of my talk will concern contributions^a to point **(ii)**, with a discussion of the phenomenological relevance of (resummed) NLO-BFKL kernels and point **(iii)**, with a discussion of traveling wave solutions of non-linear QCD equations, as being deeply related to geometric scaling and the transition to saturation.

^aWe shall leave the point **(i)** outside of the scope of the present conference, despite some recent developments related to the AdS/CFT correspondence¹⁰ and the “BFKL treatment” of the 4-Supersymmetrical gauge field theory¹¹.

2. “Instability towards DGLAP”

The “instability” of the BFKL equation’s solution w.r.t. the renormalization group evolution is well-known¹². Indeed, the first correction to the leading-log $1/x$ approximation of the BFKL kernel³ is large enough to apparently endanger the whole BFKL approach. It was soon realized that a large part of the problem was due to the appearance of singularities which contradict the renormalization group properties. Hence requiring an harmonization between the next leading log BFKL calculations and the renormalization group requirements through higher orders’ resummation leads^{4,5} to a possible way out of the problem.

Let us focus¹³ on the impact of these developments on the proton structure functions and recall the parametrization of the proton structure functions in the (LO) BFKL approximation¹⁴:

$$F_i = \int \frac{d\gamma}{2i\pi} \left(\frac{Q^2}{Q_0^2} \right)^\gamma x_{Bj}^{-\frac{\alpha_s N_c}{\pi} \chi_{L0}(\gamma)} h_i(\gamma) \eta(\gamma) , \quad (1)$$

where F_i denotes respectively F_T , F_L , G (resp. transverse, longitudinal and gluon) structure functions and h_i are the known perturbative couplings to the photon ($h_G=1$ for the gluon structure function), usually called “impact factors”¹⁵. χ_{L0} is the the LO-BFKL kernel, α_s the (fixed) coupling constant and $\omega(\gamma)$ an (unknown but factorizable¹⁵) non-perturbative coupling to the proton. Mellin-transforming (1) in $j-1 \equiv \omega$ space, one easily finds

$$\tilde{F}_i = \int \frac{d\gamma}{2i\pi} \left(\frac{Q^2}{Q_0^2} \right)^\gamma \frac{1}{\omega - \frac{\alpha_s N_c}{\pi} \chi_{L0}(\gamma)} h_i(\gamma) \eta(\gamma) , \quad (2)$$

and, taking the pole contribution, one has the important relation

$$\omega = \frac{\alpha_s N_c}{\pi} \chi_{L0}(\gamma_k(\omega)) . \quad (3)$$

Let us try and find the equivalent relation at NLO. At (resummed) next-to-leading order, one can similarly write^b

$$\tilde{F}_i = \int \frac{d\gamma}{2i\pi} \left(\frac{Q^2}{\Lambda_{QCD}^2} \right)^\gamma e^{-\frac{x(\gamma,\omega)}{b\omega}} h_i(\gamma, \omega) \eta(\gamma, \omega) , \quad (4)$$

^bEq.(4) is already an approximation of the (still) unknown complete (resummed) NLO-BFKL formula, since the photon and proton impact factors are not yet known at NLO. However, one expects Eq.(4) to be a phenomenologically valid approximation containing the information on the NLO kernel.




Figure 2. $\bar{\gamma}(\omega, Q^2)$: The “effective” F_2 anomalous dimension. $\bar{\gamma}(\omega, Q^2)$ has been evaluated from four known different parametrizations. They are all compatible in the range $.3 < \omega < 1, 5 < Q^2 < 100 \text{ GeV}^2$, where we restrict our analysis.


where, by construction

$$\frac{\partial}{\partial \gamma} X(\gamma, \omega) \equiv \chi_{NLO}(\gamma, \omega) . \quad (5)$$

The function $X(\gamma, \omega)$ appears in the solution of the Green function derived^c from the renormalization-group improved small- x_{Bj} equation⁴, $\chi_{NLO}(\gamma, \omega)$ is a resummed NLO-BFKL kernel and

$$\frac{N_c}{\pi} \alpha_s(Q^2) = [b \ln(Q^2/\Lambda_{QCD}^2)]^{-1} , \quad (6)$$

^cThe second variable of $X(\gamma, \omega)$ in (5) corresponds to the choice of a reference scale $\mu \rightarrow \omega \equiv j-1$ dictated by the treatment of the Green function fluctuations near the saddle-point⁴.

Figure 3. *Test of Relation ii): LO-BFKL.*


with $b = 11 - 2/3 N_f/N_c$.

At large enough Q^2/Λ_{QCD}^2 , one can use the saddle-point approximation in γ to evaluate (4). Assuming that the perturbative impact factors and the non-perturbative function η do not vary much^d, the saddle-point condition reads

$$\omega \sim \frac{\chi_{NLO}(\bar{\gamma}, \omega)}{b \ln(Q^2/\Lambda_{QCD}^2)} = \frac{N_c \alpha_s(Q^2)}{\pi} \chi_{NLO}(\bar{\gamma}, \omega), \quad (7)$$

where $\bar{\gamma} \equiv \bar{\gamma}(\omega, Q^2)$ is the saddle-point value.

^dWe do not take into account modifications *e.g.* coming from powers of γ in the prefactors which may shift the saddle point⁴. We thus assume a smoothness of the structure function integrand around the saddle-point in agreement with the phenomenology¹³.

Figure 4. *Test of Relation ii): NLO-BFKL.*

Inserting the saddle point defined by (7) in formula (4), one obtains a set of conditions to be fulfilled at (resummed) NLO level as follows:

- i) The Mellin transform $\tilde{F}_2 \equiv \tilde{F}_T + \tilde{F}_L$ defines:

$$\frac{\partial}{\partial \ln(Q^2)} \ln \tilde{F}_2(\omega, Q^2) \sim \tilde{\gamma}(\omega, Q^2). \quad (8)$$

- ii) $\tilde{\gamma}$ verifies

$$\chi_{NLO}(\tilde{\gamma}) \equiv \frac{\pi \omega}{\alpha_s(Q^2) N_c}, \quad (9)$$

where χ_{NLO} is a resummed NLO-BFKL kernel.

- iii) The gluon structure function (one may also choose the observable

F_L) verifies, *via* Mellin transform:

$$\ln(\tilde{G}(\omega, Q^2)) = \ln(\tilde{F}_2(\omega, Q^2)) - \bar{\gamma} \ln[h_T(\bar{\gamma}) + h_L(\bar{\gamma})] . \quad (10)$$

We test¹³ the properties **i)-iii)** using NLO kernels proposed in⁴, and compared with the LO kernel condition (3).

As an example an extraction of an “effective” F_2 anomalous dimension **i)**, see Fig. (2), is performed¹³ using different parametrisations in a range of ω verifying the stability with respect to cuts on unknown (smallest) or irrelevant (large) x_{Bj} . The comparison of the property **ii)** to the LO BFKL kernel is displayed in Fig.(2) and the one with a resummed NLO-BFKL kernels (*cf.* Scheme⁴ 3) in Fig.(2). As is clearly seen on the figures the Mellin-transform analysis disfavors the BFKL-LO kernel, while it is qualitatively compatible with the resummed BFKL-NLO kernel. The remaining discrepancies at NLO could be attributed to finite NNLO corrections to the kernel or to still unknown NLO contributions to the impact factors¹⁶. A systematic study of the proposed NLO kernels is thus made possible using the method^e.

3. “Saturation instability”

As well-known, the BFKL evolution (even including next-leading contributions) leads to a multiplication of partons with non-vanishing size and thus inevitably leads to a dense medium . This may be called the “Saturation Instability” of the BFKL evolution.

The back-reaction of parton saturation on the BFKL equation has been originally⁷ described by adding a non-linear damping term. More recently, the evolution equation to saturation have been theoretically derived in the case of scattering on a “large nucleus”, *e.g.* when the development of the parton cascade is tested by uncorrelated probes^{8,9}.

In the following we will focus on the solutions of the Balitsky-Kovchegov (BK) equations⁹ where one consider the evolution within the QCD dipole Hilbert space¹⁷. To be specific let us consider $N(Y, x_{01})$, the dipole forward scattering amplitude and define

$$\mathbb{N}(Y, k) = \int_0^\infty \frac{dx_{01}}{x_{01}} J_0(kx_{01}) N(Y, x_{01}) . \quad (11)$$

Within suitable approximations (large N_c , summation of fan diagrams,

^eSimilarly, relation **iii)** can be looked at using the gluon structure function parametriza-




Figure 5. *Typical traveling wave solution.* The function $u(t, x)$ is represented for three different times. The wave front connecting the regions $u = 1$ and $u = 0$ travels from the left to the right as t increases. That illustrates how the “strong absorption” or saturated phase region invades the “transparency” region.

spatial homogeneity) and starting from the Balitsky-Kovchegov equation⁹, it has been shown that this quantity obeys the nonlinear evolution equation

$$\partial_Y \mathbb{N} = \bar{\alpha} \chi(-\partial_L) \mathbb{N} - \bar{\alpha} \mathbb{N}^2, \quad (12)$$

where $\bar{\alpha} = \alpha_s N_c / \pi$, $\chi(\gamma) = 2\psi(1) - \psi(\gamma) - \psi(1-\gamma)$ is the characteristic function of the BFKL kernel¹, $L = \log(k^2/k_0^2)$ and k_0 is some fixed low momentum scale. It is well-known that the BFKL kernel can be expanded to second order around $\gamma = \frac{1}{2}$, if one sticks to the kinematical regime $8\bar{\alpha}Y \gg L$. We expect this commonly used approximation to remain valid for the full nonlinear equation. The latter boils down to a parabolic nonlinear partial derivative equation:

$$\partial_Y \mathbb{N} = \bar{\alpha} \bar{\chi}(-\partial_L) \mathbb{N} - \bar{\alpha} \mathbb{N}^2, \quad (13)$$

with

$$\bar{\chi}(-\partial_L) = \chi\left(\frac{1}{2}\right) + \frac{\chi''\left(\frac{1}{2}\right)}{2} (\partial_L + \frac{1}{2})^2. \quad (14)$$

The key point of our recent approach¹⁸ is to remark that the structure of Eq.(13) is identical (for fixed α) to a mathematical and physical archetype of

tions. However assumptions on the perturbative make the conclusions more qualitative or indicating some discrepancies to be solved at NLO.

non-linear evolution equation for which useful tools can be applied, namely the Fisher and Kolmogorov-Petrovsky-Piscounov (KPP) equation¹⁹ for a function u :

$$\partial_t u(t, x) = \partial_x^2 u(t, x) + u(t, x)(1 - u(t, x)) , \quad (15)$$

which is directly related¹⁸ to \mathbb{N} . The equation can be generalized to many physical situations, including running α .




Figure 6. *Evolution of the reduced front profile.* Fixed coupling: left; Running coupling: right. The reduced front profile $(k^2/Q_s^2)^{\gamma_c} \mathbb{N}(k/Q_s(Y), Y)$ is plotted against $\log(k^2/Q_s^2)$ for different rapidities. The various lines correspond to rapidities from 2 (lower curves, full line) up to 10 (upper curves, dotted). Note the similarity of the wave fronts, but the quicker time evolution (in \sqrt{t}) for fixed coupling, by contrast with the slow time evolution (in $t^{1/3}$) for the running coupling case.

Our main results are the following. The well-known geometric scaling property²⁰ is obtained for the solution of the non-linear equation (13) for the gluon amplitude at large energy. In our notation, the geometric scaling property reads

$$\mathbb{N}(Y, k) = \mathbb{N}\left(\frac{k^2}{Q_s^2(Y)}\right) , \quad (16)$$

where $Q_s^2(Y)$ is the saturation scale. We prove that geometric scaling is directly related to the existence of traveling wave solutions of the KPP equation¹⁹ at large times. This means that there exists a function of one variable w such that

$$u(t, x) \underset{t \rightarrow +\infty}{\sim} w(x - m(t)) \quad (17)$$

uniformly in x . Such a solution is depicted on Fig.(5). The function $m(t)$ depends on the initial condition. For the QCD case¹⁸, one has to consider

$$m(t) = 2t - \frac{3}{2} \log t + \mathcal{O}(1) . \quad (18)$$

When transcribed in the appropriate physical variables, this mathematical result, implies directly the known geometric scaling properties¹⁸. It is interesting to note how the traveling wave solutions provide a particularly striking mathematical realization of an “instability” as depicted in Fig.(5), when a stable fixed point (strong absorption) “invades” an unstable one (transparency).

This mathematical analysis can be extended¹⁸ to the study of the transition towards geometrical scaling , *i.e.* the formation of the front wave as a function of time, both for fixed and running α , see Fig.(6).

4. Conclusion

In the present contribution, we have discussed some aspects of the “instabilities” of the BFKL equations, *i.e.*:

- **i)** *Instability towards the non-perturbative regime*
- **ii)** *Instability towards the renormalization group regime*
- **iii)** *Instability towards the high density (saturation) regime*

At first sight, these instabilities could have appeared as drawbacks of the whole approach. On the very contrary, we have seen that the extensions of BFKL equation raised up by the treatment of “instabilities” appear to be the building blocks of most interesting recent developments towards a better understanding of QCD dynamics. As an example, I chose to present some personal recent contributions to this discussion, which are far from giving an idea of the whole extent of the works^f which attack the problem nowadays.

As a brief outlook, let us mention:

About Point **(i)**, not discussed here, let us mention the formal but informative discussion on the $N = 4$ supersymmetric QCD field theory and the AdS/CFT correspondence¹¹.

Point **(ii)**: It is the subject of a developing activity which will allow to master the rather high technicality of the BFKL-NLO calculations and

^fI mentioned quite a few of them in the reference list but I want to apologize for the authors and studies which I may have forgotten in this necessarily shortened review.

thus to penetrate the subtle aspects of the compatibility between BFKL and DGLAP evolution equations.

Point (iii): Saturation with both its phenomenological and theoretical aspects will certainly retain the attention of the Particle Physics community. The challenge here is the quest for a new phase of intense QCD fields and the understanding of its dynamical properties.

Acknowledgments

I want to warmly thank my collaborators in the work which has been discussed here: Stéphane Munier, Christophe Royon, Laurent Schoeffel and many colleagues with whom I had fruitful discussions, including those taking place in the charming decor of Ringberg Castle, in front of the Bavarian Alps.

References

1. L.N.Lipatov, *Sov.J.Nucl.Phys.* **23** (1976) 642; V.S.Fadin, E.A.Kuraev and L.N.Lipatov, *Phys. Lett.* **B60** (1975) 50; E.A.Kuraev, L.N.Lipatov and V.S.Fadin, *Sov.Phys.JETP* **44** (1976) 45, **45** (1977) 199; I.I.Balitsky and L.N.Lipatov, *Sov.J.Nucl.Phys.* **28** (1978) 822.
2. J.Bartels, H.Lotter, M. Vogt, *Phys. Lett.* **B373** (1996) 210.
3. V.S. Fadin and L.N. Lipatov, *Phys. Lett.* **B429** (1998) 107; M.Ciafaloni, *Phys. Lett.* **B429** (1998) 363; M. Ciafaloni and G. Camici, *Phys. Lett.* **B430** (1998) 349.
4. G.P. Salam,, *JHEP 9807* (1998) 019. M. Ciafaloni, D. Colferai, G.P. Salam,, *Phys.Rev.* **D60** 114036, , *JHEP 9910* (1999) 017; M. Ciafaloni, D. Colferai, G.P. Salam, A.M. Stasto,, *Phys. Lett.* **B541** (2002) 314.
5. S.J. Brodsky, V.S. Fadin, V.T. Kim, L.N. Lipatov, G.B. Pivovarov, *JETP Lett.* **70** (1999) 105.
6. G.Altarelli and G.Parisi, *Nucl. Phys.* **B106** 18C (1977) 298. V.N.Gribov and L.N.Lipatov, *Sov.J.Nucl.Phys.* (1972) 438 and 675. Yu.L.Dokshitzer, *Sov.Phys. JETP.* **46** (1977) 641. For a review, see *e.g.* Yu.L. Dokshitzer, V.A. Khoze, A.H. Mueller, S.I. Troyan, *Basics of Perturbative QCD* .
7. L.V. Gribov, E.M. Levin and M.G. Ryskin, *Phys.Rep.* **100**, 1 (1983); A.H. Mueller, J. Qiu, *Nucl. Phys.* **B268**,(1986) 427.
8. L. McLerran and R. Venugopalan, *Phys. Rev.* **D49**, 2233 (1994); *ibid.*, 3352 (1994); *ibid.*, **D50**, 2225 (1994); A. Kovner, L. McLerran and H. Weigert, *Phys. Rev.* **D52**, 6231 (1995) ; *ibid.*, 3809 (1995); R. Venugopalan, *Acta Phys.Polon.* **B30**, 3731 (1999); E. Iancu, A. Leonidov, and L. McLerran, *Nucl. Phys.* **A692**, 583 (2001); *idem*, *Phys. Lett.* **B510**, 133 (2001); E. Iancu and L. McLerran, *Phys. Lett.* **B510**, 145 (2001); E. Ferreiro, E. Iancu, A. Leonidov

- and L. McLerran, *Nucl. Phys.* **A703**, 489 (2002); H. Weigert, *Nucl. Phys.* **A703**, 823 (2002).
9. I. Balitsky, *Nucl. Phys.* **B463**, 99 (1996); Y. V. Kovchegov, *Phys. Rev.* **D60**, 034008 (1999); *ibid.*, **D61**, 074018 (2000).
 10. J. Maldacena, *Adv.Theor.Math.Phys.* **2** (1998) 231; S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Phys. Lett.* **B428** (1998) 105; E. Witten, *Adv. Theor. Math. Phys.* **2** (1998) 253; O. Aharony, S.S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, *Phys. Rep.* **323** (2000) 183.
 11. R.A. Janik and R. Peschanski, *Nucl. Phys.* **B565** (2000) 193; *Nucl. Phys.* **B625** (2002) 279; R.A. Janik, *Phys. Lett.* **B500** (2001) 118; J. Polchinski and M.J. Strassler, *Phys. Rev. Lett.* **88** (2002) 031601. A.V. Kotikov, L.N. Lipatov, V.N. Velizhanin *Phys. Lett.* **B557** (2003).
 12. See, for instance, Kopeliovitch B.Z., Peschanski R.: *Working Group report on Diffraction Highlights of the Theory*, Proceedings of the “6th International Workshop on Deep Inelastic Scattering and QCD (DIS 98) COREMANS G., ROOSEN R., eds. pp. 858-862, (World Scientific, 1998) Brussels, Belgium, 1998.
 13. R.Peschanski, C.Royon and L.Schoeffel, to appear. See a preliminary discussion in R.Peschanski, *Acta Phys. Pol.* **34** (2003) 3001; L.Schoeffel, contribution to the ‘11th International Workshop on Deep Inelastic Scattering and QCD (DIS 03), Proceedings to appear.
 14. H Navelet, R.Peschanski, Ch. Royon, S.Wallon, *Phys. Lett.* **B385** (1996) 357. S.Munier, R.Peschanski, *Nucl. Phys.* **B524** (1998) 377.
 15. S. Catani, M. Ciafaloni, F. Hautmann, *Nucl.Phys.* **B366** (1991) 135; S. Collins, R.K. Ellis, *Nucl.Phys.* **B360** (1991) 3; E.M. Levin, M.G. Ryskin, Yu.M. Shabelski, A.G. Shuvaev, *Sov.J.Nucl.Phys.* **B53** (1991) 657.
 16. J. Bartels, D. Colferai, S. Gieseke, A. Kyrleis, *Phys.Rev.* **D66** 094017, and references therein.
 17. A.H. Mueller, *Nucl.Phys.* **B415** (1994) 373; A.H. Mueller and B. Patel, *Nucl.Phys.* **B425** (1994) 471; A.H. Mueller, *Nucl.Phys.* **B437** (1995) 107.
 18. S. Munier and R. Peschanski, “*Geometric scaling as traveling waves*”, to appear in *Phys. Rev. Lett.*, [arXiv:hep-ph/0309177]; “*Traveling wave fronts and the transition to saturation*”, [arXiv: hep-ph/0310357].
 19. R. A. Fisher, *Ann. Eugenics* **7** (1937) 355; A. Kolmogorov, I. Petrovsky, and N. Piscounov, *Moscow Univ. Bull. Math.* **A1** (1937) 1. For Mathematical properties : M. Bramson, *Memoirs of the American Mathematical Society* **44** (1983) 285. For Physical Applications, see B. Derrida and H. Spohn, E. Brunet and B. Derrida, *Phys. Rev.* **E56**(1997) 2597 ; For a recent general review, Wim van Saarloos, *Phys. Rep.* **386** (2003) 29.
 20. A. M. Stašo, K. Golec-Biernat, and J. Kwiecinski, *Phys. Rev. Lett.* **86** (2001) 596 .